

CALBO

LIQUID Calcium Boron

A Complexed Formulation for the Correction and Prevention of Calcium & Boron Deficiencies

☐

GUARANTEED ANALYSIS

Total
Calcium (Ca).....6.0%
Derived from Calcium Sulfate
Boron (B)2.0%
Derived from Sodium Borate
Wt per gal: 10.45 lb. at 68°F Grams per liter: 1252 g @20°C

☐

GENERAL INFORMATION

Calbo is designed as a foliar nutritional spray to supply Calcium and Boron in a convenient, soluble form. CaB is completely available and absorbed on the leaf surface or by the root system because of the natural complexing properties of citric acid.

COMPATIBILITY

Calbo is compatible with most pesticides. In mixing, always add this product to the tank last. A compatibility test is recommended if the desired combination has not been used in the past.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all recommended governmental procedures for disposal of pesticide spray solutions. Rinse container, empty rinsate and spray tank. Contact county agricultural commissioner for container disposal information.

CONDITIONS OF SALE

The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, the manufacturer does not make nor does it authorize any agent or representative to make any warranty or representation, expressed or implied, concerning this material or the use thereof except in conformity with the statements on the label. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions. If these conditions are unacceptable to the buyer, he should immediately return the unopened product to the seller.

Net Contents: .☐

Net Weight:

Lot #:

DIRECTIONS

For best results, spray when the crop is in an active growing state after irrigation or natural rain fall. Spray early in the morning or late afternoon. Midday sprays may not be effective because of excessive moisture evaporation.

CAUTION:

Excessive amounts of Boron may cause damage to susceptible crops. Determine plant sensitivity prior to use. Contact your local agricultural authority for information on specific crops.

Recommendations for Use: Calbo may be used on the following crops: Alfalfa, Almonds, Avocados, Apples, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Citrus, Clover, Corn, Cotton, Grapes, Lettuce, Milo, Melons, Nectarines, Rice, pears, Peaches, Pecans, Peppers, Plums, Prunes, Potatoes, Peanuts, Sorghum, Sugar Betts, Sweet Corn, Sugarcane, Strawberries, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Vegetables:

Apply 1 pint per acre per week in the normal spray program. Increase the rate to 2 pints per acre per week prior to the time that the deficiency usually shows up.

Tree Crops:

Apply 1 to 2 pints per 100 gallons of spray. Start two to three weeks after full bloom and continue at two-week intervals. Trees must be wet thoroughly. For best results do not spray in the heat of the day.

SHAKE BEFORE USE

Specimen Label

Peanuts: Apply 1 gallon per acre per application. Repeat 3 times starting at early bloom. **Cotton:** Apply a total of 1 gallon per acre with two applications of 1/2 gallon each or four treatments of 1/4 gallon per acre at 10 to 14 day intervals starting at early bloom. **Soybeans:** Apply 1/2 gallon per acre at early pod set and an additional application of 1/2 gallon per acre in 14 days.
Note: Do not exceed 1 gallon per acre per application.

Foliar Applications:

Conventional Sprayers: Use a minimum of 20 gallons of water per acre.

Low Volume Sprayers: Use 5 gallons of water per acre. If less water is used, slight burning of the foliage may occur.

Aerial Applications: Do not exceed 1 quart per gallon of water

PRECAUTIONARY STATEMENTS

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid skin contact. Causes eye and skin irritations. In case of contact, wash the affected areas with water immediately and continue washing for fifteen minutes. Seek medical attention if irritation persists.

STATEMENT OF PRACTICAL TREATMENT

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area and seek medical attention, if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

SHAKE WELL BEFORE USE

MANUFACTURED BY:

CUSTOM AGRICULTURAL FORMULATORS
P.O. BOX 26104 - FRESNO, CALIFORNIA 93729

☐ ☐ ☐ ☐